

Latest Vaccination Chart and Schedule in India

Age	Content Tag	Vaccines	Mandatory and Optional Vaccines	Important Notes
Birth	BCG	BCG (Bacille Calmette Guerin)	BCG vaccine - for tuberculosis is Mandatory	
	HB 1	Hepatitis B	Mandatory	
	OPV 0	Polio	Mandatory	
6 weeks	HB 2	Hepatitis B	Mandatory	
	IPV 1	Polio	Mandatory	In case IPV is not available or feasible, the child should be offered bOPV (3 doses). In such cases, give two fractional doses of IPV at 6 week and 14 week
	DTP 1	DTwP/DTaP (Diphtheria and Tetanus toxoids and Whole-cell Pertussis vaccine)	DTP vaccine - for diphtheria, tetanus, and pertussis (whooping cough) is Mandatory	
	HiB 1	HiB (Haemophilus Influenza B)	Haemophilus Influenza B Vaccine is Mandatory	
	PCV 1	Pneumococcal conjugate vaccine	Optional	
	Rota 1	Rotavirus	Mandatory	

10 weeks	HB 3	Hepatitis B		
	IPV 2	Polio		
	DTP 2	DTwP/DTaP		
	HiB 2	HiB		
	PCV 2	Pneumococcal		
	Rota 2	Rotavirus		
14 weeks	HB 4	Hepatitis B		Fourth dose of Hepatitis B permissible for combination vaccines only
	IPV 3	Polio		
	DTP 3	DTwP/DTaP		
	HiB 3	HiB		
	PCV 3	Pneumococcal		
	Rota 3	Rotavirus		Third dose not required for RV1. Catch-up up to 1 year of age in UIP schedule
6 months	TCV	Typhoid Conjugate Vaccine	Typhoid Conjugate Vaccine is Mandatory	
	Influenza	Influenza	Influenza (flu vaccine) is Optional	Begin influenza vaccination after 6 months of age, about 2-4 weeks before season; give 2 doses at the interval of 4 weeks during first year and then single dose yearly till 5 years of age
9 months	MMR	MMR 1 (Measles, Mumps and	MMR vaccine for measles, mumps and	

		Rubella)	rubella is Mandatory	
	MCV 1	Meningococcal	Meningococcal meningitis is Optional	Meningococcal Vaccine (MCV) 9 months through 23 months: 2 doses, at least 3 months apart; 2 years through 55 years; single dose only
12 months	Hep A1		Hepatitis A vaccine is Mandatory	
	MCV 2	Meningococcal		
	JE 1	JE	Japanese encephalitis vaccine is given in high risk areas only, or in case of travel to high-risk areas – So it is Optional	For individuals living in endemic areas & for travellers to JE endemic areas provided their expected stay is for a minimum period of 4 weeks
	Cholera	Cholera	Cholera vaccine is given in high risk areas only, or in case of travel to high-risk areas. So it is Optional	Two doses two weeks apart for >1 year old; for individuals living in high endemic areas and travelling to areas where risk of transmission is very high
13 months	JE 2	JE		
15 months	PCV B1	Pneumococcal		
	MMR	MMR 2		
	Varicella	Varicella 1	Chickenpox (varicella) is Optional	

		Cholera 1 & 2		
16-18 months	IPV B1	Polio		b-OPV, if IPV booster (standalone or combination) not feasible
	DTwP/ DTaP	DTP B1		
	HiB	HiB B1		
	Hep A2	Hepatitis A		Live attenuated Hepatitis A vaccine: single dose only
2-3 years	MCV	Meningococcal		
4-5 years	DTwP/ DTaP	DTP B2		
	MMR	MMR3/ MMRV		
	Varicella	Varicella 2		
9-14 years	PCV	Pneumococcal		
	Tdap	Tdap		
	HPV	Human Papilloma Virus 1 & 2		2 doses at 6 months interval 9-14 years age; 3 doses (at 0, 1-2 & 6 months) 15 years or older and immunocompromised
15-18 years	Tdap	Td		
	HP 1,2,3	HPV 1,2,3		